



International Journal of Advance Research in Nursing

Volume 4; Issue 1; Jan-June 2021; Page No. 123-129

Received: 25-11-2020
Accepted: 29-12-2020

Indexed Journal
Peer Reviewed Journal

Perceptions of Kuwaiti Heads of Education and Health Departments Towards School Nurses in Tackling Obesity Among School Children

¹Dr. Mohammad Alresheedy, ²Dr. Lina Shahwan-AKL, ³Budoor Al Hajeri, ⁴Dr. Ali Alhaiti and ⁵Dr. Mishal Aldaihani

¹ Lecturer, Kuwait Nursing Institute, Public Authority for Applied Education and Training, Kuwait

² Director of Studies at Institute of Health and management, Kuwait

³ Trainer at public authority of apply training and education, Kuwait

⁴ Education consultant at King Fahad Medical City, Kuwait

⁵ Director of scholarship and Training in medical cities- Ministry of interior, Kuwait

Abstract

School nurses have been identified to be an important facilitator in helping students achieve a healthy weight. This exploratory qualitative study was conducted to examine the perceptions of officials from the Ministries of Education and Health towards school nurse in tackling obesity among children. The findings of this study show that most of the participants agreed that school nurses do not play a role in either promoting children's health or the prevention of disease. This is not in line with several studies that found that the activities of school nurses, such as delivering health education and taking body weight measurements.

Keywords: Obesity, Kuwait, school children, school nurse, perception

Introduction

Globally, the prevalence of overweight and obese children and adolescents has reached epidemic proportions (WHO, 2020) [50]. The data presented on the incidence of overweight and obesity in developed and developing countries indicate the seriousness of this issue. For instance, in developed countries, within the European region, a study conducted in 15 European countries (Spain, Italy, Belgium, Greece, Czech Republic, Hungary, Ireland, Latvia, Lithuania, Malta, Norway, Cyprus, Portugal, Slovenia and the Republic of Macedonia) presents alarming statistics showing that the rate of overweight ranged between 18% and 50% among girls and between 18% and 57% among boys (Wijnhoven *et al.*, 2014) [49].

In developing countries, the prevalence of obesity among children and adolescents was found to be 22.1 % in Brazil, 41.8 % in Mexico, 19.3% in Argentina and 22.0% in India (Gupta, Shah, Nayyar & Misra, 2013) [18]. A review in the Eastern Mediterranean Region (EMR) showed that between 7% and 45% of school children were overweight or obese (Musaiger, 2011) [31]. Several studies conducted in Kuwait show that overweight and obesity are indeed prevalent in this country, with estimates suggesting that one third of children and adolescents are overweight or obese (El-Bayoumy, Shady & Lotfy, 2009; Musaiger *et al.*, 2012; Ng *et al.*, 2011; Zaghoul, 2013) [15, 32, 53]. Research studies have shown that individuals who were overweight or obese in

childhood are more likely to have adverse health consequences during adulthood (Reilly & Kelly, 2011; WHO, 2020) [40, 50]. Several studies have also identified that obese children are more likely than healthy-weight children to become obese adults (Biro & Wien, 2010; Center for Disease Control, 2013) [7, 11].

Overweight and obesity are preventable conditions that have been identified as primary contributing factors in chronic disease (WHO, 2020) [50]. In fact, weight gain has been identified as contributing to more than 34% of mortality in people younger than 60 years of age in most Arab-speaking countries (Rahim *et al.*, 2014) [39]. Dattilo *et al.* (2012) [13] suggest that interventions must begin in the early years and focus on children who are at risk of obesity. This period of life is recognised as the best time to instil behaviours associated with a healthy lifestyle, such as food preferences and physical activity that may then extend into adulthood (Campbell *et al.*, 2014) [10]. Therefore, it can be anticipated that without early and effective interventions obesity will continue to be a public health concern in many countries.

A review of the current literature did not identify any Kuwaiti studies that examined the role of the school nurses in addressing child obesity. This is a significant gap in the literature that the current study attempts to address. A number of studies indicate that school nurses play a significant role in promoting a healthy weight for school children (Wright, Giger, Norris and Suro, 2013; National

Association of School Nurses, 2013) [51] Therefore, this study was conducted to examine the perceptions of officials from the Ministries of Education and Health towards school nurse in tackling obesity among children. Hence, this study aims to inform the development of new public health policies to respond to this serious health problem.

Method
Study Design

A qualitative method, one-to-one interviews were used to explore the role of school nurses towards obesity among school children in the state of Kuwait. with officials from the Ministries of Education and Health (six officials from the Health Department and six officials from the Education Department).

Interviews data collection

A total of 12 interviews were conducted with six government officials from the Health Department and six government officials from the Education Department in each province of the state of Kuwait. The reason for inviting officials from the departments in each province was to achieve representativeness of the two areas of responsibility across the six provinces.

Before the interviews were conducted, invitations were sent to representative officials in each province, which included a plain language letter explaining the study that provided the name and the contact details of the researcher and the supervisors. The voluntary nature of participation was explained in this letter and it was accompanied by a consent form that was signed prior to the involvement in the face-to-face interviews. Participants they were asked to indicate whether Kuwaiti schools employ school nurses and what was the attitude toward school nurses in terms of their role in contributing to contain obesity and excess weight in the schools.

Qualitative data analyses

All interviews were audiotaped and were in the Arabic language, because all the participants were Arabic speakers

from Kuwait and most did not speak English. The interviews were recorded using an MP3 device in addition to a mobile device in case there was a malfunction in the recording device, hence there was a backup. The data from the interviews was transcribed in Arabic to ensure that no data was missing and then all of this data was translated into English by a professional translation office in Kuwait. The qualitative data was analysed using content analysis. The researcher identified, coded and categorised the patterns that emerged from the data. The common statements were used to generate themes that represented the perceptions of the group. These statements were grouped, or clustered, into thematic categories. These thematic categories are presented as part of the findings along with the frequency of occurrence among the participants. In addition, textual data offered as verbatim examples from the participants discussions were included to highlight the key common responses and to clarify these concepts. Relevant data were coded by researcher throughout the document following an inductive process, which allowed for emergent categories. NVivo 10® qualitative analysis software was used during the process of coding and the development of themes. NVivo software provided an organised workspace to conduct the classification and sorting of data and the tracking of the frequency of occurrences across the data sources (Ritchie *et al.* 2013) [41].

Finally, a comprehensive review and interpretation of the data provided the conclusions of the analysis, which were revealed in the form of several overarching themes representing the perceptions of the individual interviews as a whole (Bazeley & Jackson, 2013) [6].

One-to-one interview results

The data obtained from the 12 individual interviews with government representatives from the ME and the MOH were analysed to reveal main thematic category, which were then further separated into sub-categories, the thematic category is presented individually and include main category: Presence and function of school nurses, which is described in Table 1 showing sub-category and highlighting the common themes in each.

Table 1: Categories, Sub-categories and Themes for Interview Data

Primary category	Sub-category	Themes
Presence and Function of School Nurses	(Q6) Presence of nurses in schools	Schools employ nurses through private company Nurses only in some schools-nursing shortage Yes, the schools have nurses
	Function/Role of school nurses	No role in obesity prevention, only first aid Need for more qualified nurses, need for training Focus on treatment, not prevention Trained to educate about and prevent obesity, but not used for prevention Important role in health education Language barriers Confront health problems and work for health awareness/early detection

Presence of school nurses

In discussing the presence of nurses in schools, the majority of participants (10 of 12) described a disparity between schools, where some schools had access to a school nurse, while others did not. This discrepancy was noted by five participants as being due to a shortage of available nurses to work in schools. The nurses that were employed by schools were described as being recruited through private company sources. Table 2 provides the full variety of responses and

associated frequencies.

Table 2: Availability of School Nurses

Response theme	Frequency among focus groups (n = 12)
Some, but not all, schools have a nurse	10
Some schools employ nurses through private company	6

Nurses only in some schools due to shortage of nurses	5
Schools have nurses	3

The participants generally agreed that nursing staff are not always available to all schools, which was attributed to a shortage in nurses. The following examples shed light on this problem:

Yes, the Ministry of Health used to provide nurses for the schools. We tried to cover all the schools; however, sometimes one nurse would have to cover three or four schools because of the shortage of nurses. Nowadays, school nurses are under the Ministry of Education and are employed through private companies. (A.2)

Yes. At first, these nurses were provided by the Ministry of Health; however, now they are provided by the Ministry of

Education, which tasks private companies with providing school nurses. The Ministry of Health is unable to provide these nurses because of the shortage in nursing staff. (C.2)

Role of nurses in schools

The perceived role of the school nurse-if present-is also critical. From the data, for schools in which a nurse was employed the role of the nurse was limited to first aid in most cases, and they played no role in prevention or education (9 of 12 participants). Despite the need for education and prevention and the presence of qualified nursing staff, the nurses were reportedly not used in this capacity. Table 3 illustrates the response variety and the frequency of occurrence in the interview data.

Table 3: Nursing Role in Schools

Response theme	Frequency among focus groups (n = 12)
No role in prevention; role is focused on treatment and first aid	9
Need for training and work program; need for more qualified nurses	3
Are trained and educated to handle role of reducing obesity, but not used in this capacity	2
Important role in health education, but not used in this capacity	2
Language barriers	2
Confront health problems and work for health awareness for students; role of early detection	2

Along with the problem of a shortage of nursing staff, the role of the nurses in schools was described by nine of the 12 participants as limited, primarily to first aid, and the school nurse was not utilised for prevention purposes:

Nurses are in the schools to provide first aid only. They have no role in addressing this problem. (F.1)

The school nurse does not have a role in this problem because the work of the nurse in Kuwaiti schools is only related to treatment and first aid. (J.1)

In Kuwait, there are roles that the school nurse is not asked to play, such as the prevention and curative role. Nurses have an important role in health education. (J.2)

However, three participants had concerns that the nurses were not qualified to fill the preventative role. For example, one participant stated:

I don't think that nurses in Kuwait have reached this stage because they are not qualified to play such a role. Currently, they only focus on treatment and not on prevention. We need courses and programs that will enable them to resolve this problem. (F.2)

Other participants felt the nurses were or could be qualified for the role and should receive training for this. This was demonstrated in the response by H.1, who stated: 'The school nurse's role is more about treatment than prevention, and this is wrong. I think that if nurses are trained well first, they could help through education and measuring weight and height.' If appropriately trained and given responsibility for prevention, school nurses could be a significant part of the solution to childhood obesity. This was expressed by participant C.1 as follows: 'If nurses were given more responsibilities, worked as a team in the schools and developed an intensive program for each school, then they could play a very important role in solving this problem.'

Discussion

A strong theme that emerged from the focus groups

demonstrated that school nurses were poorly utilised in Kuwaiti schools. All the participants reported that school nurses were limited to providing first aid rather than capitalising on the potential of nurses to serve an important role in health promotion and disease prevention. Similar findings from a qualitative study conducted in the US indicates that parents and teachers perceived that the role of the school nurse was to administer medications and provide first aid (Maughan & Adams, 2011). Similarly, studies on US parents (N = 369) indicated that the majority of parents (71%) perceived that the role of the school nurse was to provide emergency care, while 31% of parents perceived that nurses had a role to play in identifying health problems or performing health screenings (Kirchofer, Telljohann, Price, Dake, & Ritchie, 2007) [20]. Likewise, a study conducted in Greece revealed that 63.9% of teachers (N = 144) thought that the role of the school nurse was to provide first aid and 36% believed that the school nurse should provide treatment and health education (Mastrogiannis *et al.*, 2013) [26].

In contrast, several studies disagreed with the results of the current study (Bonde, Bentsen, & Hindhede, 2014; Wright, Giger, Norris, & Suro, 2013; Gonzaga *et al.*, 2014; Leibold, 2009; Mehrley & Leibold, 2011; Pbert *et al.*, 2013; Tucker & Lanningham-Foster, 2015; Wright, Giger, Norris, & Suro 2013) [8, 16, 51, 24, 28, 36, 48]. These studies indicate that school nurses can play a critical role in promoting children's health. Kubik and Lee (2013) [22] identified that school nurses are highly skilled members of staff who have been poorly utilised in the fight against obesity. The findings of a systematic review on the role and influence of nurses in US elementary schools highlighted many significant activities of school nurses that can be described as disease prevention or health promotion, including screening students' vision and BMI, educating students and school staff, and tracking student immunisations (Lineberry & Ickes, 2014) [25].

Pender, Murdaugh and Parsons (2011) [37] conclude that nurses can work with individuals, families and communities to create the optimal conditions for the expression of a high level of wellbeing.

Moreover, evidence provided in the literature indicates that school nurses can contribute to the improvement of adolescents' dietary habits (Pbert *et al.*, 2013) [36], increase children's physical activity (Leibold, 2009; Tucker & Lanningham-Foster, 2015; Wright *et al.*, 2013) [24, 48, 51], decrease sedentary activity (Wright *et al.*, 2013) [51] and help overweight children and adolescents adopt healthy habits (Gonzaga, Araujo, Cavalcante, Lima, & Galvao, 2014) [16]. O'Brien (2012) [16] identified that each point increase in school nurses' activities was associated with a 0.16% reduction in obesity in school students. Soto and White (2010) [43] point out that the first step toward obesity intervention starts with screening BMI and sending parents a notification. In fact, several studies' findings indicate that school nurses are screening children's body weight in their daily practice (Krause-Parello & Samms, 2009; Kubik, Story, & Davey, 2007; Lineberry & Ickes, 2014; Stalter, Chaudry, & Polivka, 2011) [21, 23, 25, 44].

Stoddard, Kubik and Skay (2008) [46] stated that school nurses are skilled staff and capable of measuring a child's BMI in a reliable and private manner. Likewise, Hendershot, Telljohann, Price, Dake and Mosca's (2008) study on school nurses (N = 2,629) found that 81% of nurses believed they should set a plan to address overweight in children, and up to 67% of the nurses believed they should measure BMI and send this information home.

Some participants in the current study suggest that school nurses can play an essential role in the prevention of disease, but this role would depend on the nurses' qualifications and responsibilities. It is true that the literature indicates that the school nurses perceived several barriers to addressing obesity, including a lack of knowledge, a lack of resources, time and support, as well as personal weight challenges (Steele *et al.*, 2011) [45]. In addition, a lack of privacy, high workloads and a lack of policy support were seen as key barriers to body weight assessment (Stalter, Chaudry, & Polivka, 2011) [44]. Moyers, Bugle and Jackson (2005) [30] found that school nurses perceived numerous barriers to taking a more active role in tackling childhood obesity, including a low level of competence and difficulty in providing counselling and a lack of parental support.

Quelly (2014) [38] summarised that school nurses need further education, policy support and professional mentoring to take part in childhood obesity intervention. Moreover, Melin and Lenner (2009) [29] found that trained school nurses with sufficient knowledge working in a collaborative team were more effective in tackling overweight and obesity in their schools. Another important barrier noted by two focus groups in the current study was a lack of communication. One participant commented that nurses do not speak Arabic, and that they faced difficulties in communicating with nurses. This is common in the Arabic world (Al-Khathami, Kojan, Aljumah, Alqahtani, & Alrwaili, 2010; Al-Jarallah, Moussa, Hakeem, & Al-Khanfar, 2009; Almutairi, McCarthy, & Gardner, 2014) [2, 3, 5], since the majority of nurses are foreign and they are not Arabic speakers. Indeed, barriers to communication are

considered a serious matter that needs to be addressed to provide a high quality of service and ensure client satisfaction.

Divi, Koss, Schmaltz and Loeb (2007) [14] mentioned that a language barrier can seriously harm patient safety-this harm ranging from moderate to death. Therefore, health providers are often obligated to provide an interpreter service (Green *et al.*, 2005; Suliman & Tadros, 2011) [17, 47]. For instance, Suliman, and Tadros (2011) [47] stated that nursing administrations in Saudi Arabia have addressed the language barrier by appointing bilingual staff as interpreters to facilitate effective communication between patients and nurses.

Finally, it was noted in the present study that there are shortages of school nurses. Not surprisingly, this is a global issue and has been reported in many countries, for example, in China (Yun, Jie, & Anli 2010) [52], in India (Saeed, 2010), in Iran (Zarea, Negarandeh, Dehghan-Nayeri, & Rezaei-Adaryani, 2009) [54] and in Saudi (Almalki, FitzGerald, & Clark, 2011) [4]. Chan, Tam, Lung, Wong and Chau (2013) [12] attribute the shortage of nurses to individual factors, including job satisfaction, demographic factors, burnout and organisational factors such as work commitment, environment, social support, culture and work demands.

Al-Kandari and Lew (2005) [1] attribute the shortage of nurses in Kuwait to the high turnover rate among foreign nurses, low production of indigenous nurses and the expansion of health care services. In contrast, Almalki, FitzGerald and Clark (2011) [4] point out that the mass media in Saudi Arabia should engage in promoting nursing as a career, and the education sector should support students financially and reduce the length of nursing training to 3 years, rather than 5 years, to encourage Saudi youth to study nursing as a specialty. Similarly, Chan, Tam, Lung, Wong and Chau (2013) [12] suggest that enhancing the work environment and increasing nurses' external rewards is needed to address the nursing shortage.

In conclusion, the findings of the current study indicate that the role of school nurses in Kuwait is under-recognised and, not surprisingly, the majority of school nurses in Kuwait are employed from private companies who look for profit rather than the quality of work. The findings of this study show that most of the participants agreed that school nurses do not play a role in either promoting children's health or the prevention of disease. However, this is not in line with the evidence provided in the literature. It seems that the shortage of nurses, language barriers and qualifications result in the underestimation of the role of school nurses.

Recommendation

The literature review provided evidence to indicate that school nurses can play an important role in designing and implementing intervention strategies to prevent obesity and promote healthy lifestyles to students (Gonzaga, Araujo, Cavalcante, Lima, & Galvao, 2014; Kubik, Story, & Davey, 2007; O'Brien, 2012) [16, 23, 35]. Therefore, it was felt that it was important to produce particular recommendations for school nurses to help nurses take part in the obesity prevention program. It is recommended that the MOH and Ministry of Education should cooperate with nurses to address all the barriers currently facing school nurses, such

as a lack of knowledge, a lack of time, a lack of resources and a lack of support. For instance, the ME has the responsibility for the education of nurses in Kuwait, so they should rethink this and provide a comprehensive education program to help nurses be more effective in terms of health education. Also, they should ensure that school clinics are adequately equipped and resourced with the necessary materials to help nurses engage in child obesity prevention programs.

Another recommendation is for the ME to recruit school nurses according to their experience and qualifications. This recommendation will ensure that nurses are able to effectively perform their duties in relation to obesity prevention in school children. In addition, the MOH has a role to play in helping nurses improve their skills and knowledge by providing intensive programs that will restore nurses' confidence in being involved in primary, secondary and tertiary intervention. The MOH also needs intervention for follow-up.

Conflict of interest

The authors declare that they have no conflict of interest.

Acknowledgments

I would like to express my deep gratitude to my supervisors, starting with Associate Professor Lina Shahwan for her expert supervision. Also, my sincere and deepest thanks to my supervisor, Dr Jennifer James, for playing an important role in this study and for always being optimistic that I also extend my special thanks to Associate Professor Phillip Maude, and I am very grateful to Dr. Linda Jones for her amazing support and guidance. Extend my thanks and gratitude to the government of the state of Kuwait for supporting me and granting me scholarship.

References

- Al-Kandari FH, Lew I. Kuwaiti high school students' perceptions of nursing as a profession: Implications for nursing education and practice. *Journal of Nursing Education* 2005;44(12):533-40.
- Al-Khathami AM, Kojan SW, Aljumah MA, Alqahtani H, Alrwaili H. The effect of nurse-patient language barrier on patients' satisfaction. *Saudi Medical Journal* 2010;31(12):1355-1358.
- Al-Jarallah KF, Moussa MAA, Hakeem SK, Al-Khanfar FK. The nursing workforce in Kuwait to the year 2020. *International Nursing Review* 2009;56(1):65-72.
- Almalki M, FitzGerald G, Clark M. The nursing profession in Saudi Arabia: An overview. *International Nursing Review* 2011;58(3):304-311.
- Almutairi AF, McCarthy A, Gardner GE. Understanding cultural competence in a multicultural nursing workforce registered nurses' experience in Saudi Arabia. *Journal of Transcultural Nursing*, 2014. 1043659614523992.
- Bazeley P, Jackson K. (Eds.). *Qualitative data analysis with NVivo*. Sage Publications Limited. Retrieved 10 October 2013 from <https://books.google.com.au/books?hl=en&lr=&id=Px8cJ3suqccC&oi=fnd&pg=PP2&dq=Qualitative+data+analysis+with+NVivo>.
- Biro FM, Wien M. Childhood obesity and adult morbidities. *American Journal of Clinical Nutrition* 2010;91(5):1499S-1505S.
- Bonde AH, Bentsen P, Hindhede AL. School nurses' experiences with motivational interviewing for preventing childhood obesity. *Journal of School Nursing* 2014;30(6):448-455.
- Boutelle KN, Cafri G, Crow SJ. Parent predictors of child weight change in family based behavioral obesity treatment. *Obesity* 2012;20(7):1539-1543.
- Campbell F, Conti G, Heckman JJ, Moon SH, Pinto R, Pungello E *et al*. Early childhood investments substantially boost adult health. *Science* 2014;343(6178):1478-1485.
- Center for Disease Control. Make a difference at your school. Chronic disease, Paper 31. 2013. Retrieved from <http://digitalcommons.hsc.unt.edu/disease/31>
- Chan ZC, Tam WS, Lung MK, Wong WY, Chau CW. A systematic literature review of nurse shortage and the intention to leave. *Journal of Nursing Management* 2013;21(4):605-613.
- Dattilo AM, Birch L, Krebs NF, Lake A, Taveras EM, Saavedra JM. Need for early interventions in the prevention of pediatric overweight: A review and upcoming directions. *Journal of Obesity* 2012;(18):123-023.
- Divi C, Koss RG, Schmaltz SP, Loeb JM. Language proficiency and adverse events in US hospitals: A pilot study. *International Journal for Quality in Health Care* 2007;19(2):60-67.
- El-Bayoumy I, Shady I, Lotfy H. Prevalence of obesity among adolescents (10 to 14 years) in Kuwait. *Asia-Pacific Journal of Public Health* 2009;3(10):117-1010.
- Gonzaga NC, Araujo TLD, Cavalcante TF, Lima FET, Galvao MTG. Nursing: Promoting the health of overweight children and adolescents in the school context. *Revista da Escola de Enfermagem da USP* 2014;48(1):153-161.
- Green AR, Ngo-Metzger Q, Legedza AT, Massagli MP, Phillips RS, Iezzoni LI. Interpreter services, language concordance, and health care quality. *Journal of General Internal Medicine* 2005;20(11):1050-1056.
- Gupta N, Shah P, Nayyar S, Misra A. Childhood obesity and the metabolic syndrome in developing countries. *The Indian Journal of Pediatrics* 2013;80(1):28-37.
- Hendershot C, Telljohann SK, Price JH, Dake JA, Mosca NW. Elementary school nurses' perceptions and practices regarding body mass index measurement in school children. *Journal of School Nursing* 2008;24(5):298-309.
- Kirchofer G, Telljohann SK, Price JH, Dake JA, Ritchie M. Elementary school parents'/guardians' perceptions of school health service personnel and the services they provide. *Journal of School Health* 2007;77(9):607-614.
- Krause-Parello CA, Samms K. The US model: The role of school nurses in New Jersey. *British Journal of School Nursing* 2009;4(6):287-292.
- Kubik MY, Lee J. Parent interest in a school-based, school nurse-led weight management program. *Journal of School Nursing* 2013. doi:1059840513485091
- Kubik MY, Story M, Davey C. Obesity prevention in

- schools: Current role and future practice of school nurses. *Preventive Medicine* 2007;44(6):504-507.
24. Leibold N. The effect of a school nurse led education intervention on blood pressure and physical activity levels in adolescents (Doctoral dissertation, College of Saint Mary) 2009. Retrieved 15 August 2015 from <http://www.csm.edu/wfdata/files/Academics/Library/InstitutionalRepository/13.pdf>
 25. Lineberry MJ, Ickes MJ. The role and impact of nurses in American elementary schools: A systematic review of the research. *Journal of School Nursing* 2014. doi: 1059840514540940
 26. Mastrogiannis D, Deltsidou A, Noula M, Poulaka MA, Gesouli-Voltyraki E, Fouka G *et al.* Exploring educationalists' views on the need for school nurses in secondary schools in Greece. *British Journal of School Nursing* 2013;8(6):303-307.
 27. Maughan E, Adams R. Educators' and parents' perception of what school nurses do the influence of school nurse/student ratios. *Journal of School Nursing* 2011;27(5):355-363.
 28. Mehrley M, Leibold N. Overweight and obesity in youth in schools: The role of the school nurse. Position statement. Revised. National Association of School Nurses (NJ1) 2011.
 29. Melin A, Lenner RA. Prevention of further weight gain in overweight school children, a pilot study. *Scandinavian Journal of Caring Sciences* 2009;23(3):498-505.
 30. Moyers P, Bugle L, Jackson E. Perceptions of school nurses regarding obesity in school-age children. *Journal of School Nursing* 2005;21(2):86-93.
 31. Musaiger AO. Overweight and obesity in eastern Mediterranean region: Prevalence and possible causes. *Journal of Obesity*, 2011.
 32. Musaiger AO, Al-Mannai M, Tayyem R, Al-Lalla O, Ali EY, Kalam F *et al.* Prevalence of overweight and obesity among adolescents in seven Arab countries: A cross-cultural study. *Journal of Obesity*, 2012, Article ID 981390, 5 pages, 2012. doi:10.1155/2012/981390
 33. National Association of School Nurses. Position statement. Overweight and obesity in youth in schools-The role of the school nurse. Silver Spring, MD: Author 2013.
 34. Ng SW, Zaghoul S, Ali HI, Harrison G, Popkin BM. The prevalence and trends of overweight, obesity and nutrition-related non-communicable diseases in the Arabian Gulf States. *Obesity Reviews* 2011;12(1):1-13.
 35. O'Brien MJF. The associations of the level of school wellness policy implementation and school nurse activities to weight status of school children in Massachusetts (Doctoral dissertation, University of Massachusetts, Boston) 2012. Retrieved 10 April 2014 from http://scholarworks.umb.edu/doctoral_dissertations/92/
 36. Pbert L, Druker S, Gapinski MA, Gellar L, Magner R, Reed G *et al.* A school nurse-delivered intervention for overweight and obese adolescents. *Journal of School Health* 2013;83(3):182-193.
 37. Pender NJ, Murdaugh CL, Parsons MA. *Health promotion in nursing practice* (6th ed.). Boston, MA: Pearson 2011.
 38. Quelly SB. Influence of perceptions on school nurse practices to prevent childhood obesity. *Journal of School Nursing* 2014;30(4):292-302.
 39. Rahim HFA, Sibai A, Khader Y, Hwalla N, Fadhil I, Alsiyabi H *et al.* Non-communicable diseases in the Arab world. *Lancet* 2014;383(9914):356-367.
 40. Reilly JJ, Kelly J. Long-term impact of overweight and obesity in childhood and adolescence on morbidity and premature mortality in adulthood: Systematic review. *International Journal of Obesity* 2011;35(7):891-898.
 41. Ritchie J, Lewis J, Nicholls CM, Ormston R. (Eds.). *Qualitative research practice: A guide for social science students and researchers*. Sage. Retrieved 2 July 2013 from [https://books.google.com.au/books?hl=en&lr=&id=EQSIAwAAQBAJ&oi=fnd&pg=PP1&dq=\).+Qualitative+research+practice:+A+guide+for+social+science+students+and+researchers.&ots=IzUPjpUs0M&sig=zHGM](https://books.google.com.au/books?hl=en&lr=&id=EQSIAwAAQBAJ&oi=fnd&pg=PP1&dq=).+Qualitative+research+practice:+A+guide+for+social+science+students+and+researchers.&ots=IzUPjpUs0M&sig=zHGM)
 42. Saeed A. Wanted: 2.4 million nurses, and that's just in India. *Bulletin of the World Health Organization* 2010;88(5):327-328.
 43. Soto C, White JH. School health initiatives and childhood obesity: BMI screening and reporting. *Policy, Politics, & Nursing Practice* 2010;11(2):108-114.
 44. Stalter AM, Chaudry RV, Polivka BJ. Regional differences as barriers to body mass index screening described by Ohio school nurses. *Journal of School Health* 2011;81(8):437-448.
 45. Steele RG, Wu YP, Jensen CD, Pankey S, Davis AM, Aylward BS. School nurses' perceived barriers to discussing weight with children and their families: A qualitative approach. *Journal of School Health* 2011;81(3):128-137.
 46. Stoddard SA, Kubik MY, Skay C. Is school-based height and weight screening of elementary students private and reliable? *Journal of School Nursing* 2008;24(1):43-48.
 47. Suliman WA, Tadros A. Nursing students coping with English as a foreign language medium of instruction. *Nurse Education Today* 2011;31(4):402-407.
 48. Tucker S, Lanningham-Foster LM. Nurse-led school-based child obesity prevention. *Journal of School Nursing* 2015. Doi: 1059840515574002
 49. Wijnhoven TM, van Raaij JM, Spinelli A, Starc G, Hassapidou M, Spiroski I *et al.* WHO European childhood obesity surveillance initiative: Body mass index and level of overweight among 6-9-year-old children from school year 2007/2008 to school year 2009/2010. *BMC Public Health* 2014;14(1):806.
 50. World Health Organization. Obesity and overweight. Fact sheet. Retrieved 110 October 2020 from <https://www.who.int/news-room/fact-sheets/detail/obesity-and-overweight>
 51. Wright K, Giger JN, Norris K, Suro Z. Impact of a nurse-directed, coordinated school health program to enhance physical activity behaviors and reduce body mass index among minority children: A parallel-group, randomized control trial. *International Journal of Nursing Studies* 2013;50(6):727-737.
 52. Yun H, Jie S, Anli J. Nursing shortage in China: State, causes, and strategy. *Nursing Outlook* 2010;58(3):122-

128.

53. Zaghoul S, Al-Hooti SN, Al-Hamad N, Al-Zenki S, Alomirah H, Alayan I *et al.* Evidence for nutrition transition in Kuwait: over-consumption of macronutrients and obesity. *Public Health Nutrition* 2013;16(04):596–607.
54. Zarea K, Negarandeh R, Dehghan-Nayeri N, Rezaei-Adaryani M. Nursing staff shortages and job satisfaction in Iran: Issues and challenges. *Nursing & Health Sciences* 2009;11(3):326-331.