



## Depression, anxiety, stress and coping strategies among patients undergoing hemodialysis

M Priyadharshini<sup>1</sup>, Liji Sara Varghese<sup>2</sup> and Dr. Hema VH<sup>3</sup>

<sup>1</sup> M.Sc. Nursing, Department of Mental Health Nursing, Dr. M.G.R Educational and Research Institute, Faculty of Nursing, Velappanchavadi, Chennai, Tamil Nadu, India

<sup>2</sup> Associate professor, Department of Mental Health Nursing, Dr. M.G.R Educational and Research Institute, Faculty of Nursing, Velappanchavadi, Chennai, Tamil Nadu, India

<sup>3</sup> Principal, Department of Medical Surgical Nursing, Dr. M.G.R Educational and Research Institute, Faculty of Nursing, Velappanchavadi, Chennai, Tamil Nadu, India

DOI: <https://doi.org/10.33545/nursing.2023.v6.i1.A.293>

### Abstract

Chronic kidney disease is an irreversible decline in renal function, which can be fatal in the absence of dialysis and transplantation. Hemodialysis causing many physical and psychological problems, such as depression, anxiety, stress and inefficiency in patient, which should be considered in health promotion intervention.

**Objectives:** To assess the level of depression, anxiety, stress among patients undergoing hemodialysis. To assess coping strategies among patients undergoing hemodialysis. To assess correlation between depression, anxiety, stress and coping strategies among patients undergoing hemodialysis. To find the association between the depression, anxiety, stress among patients undergoing hemodialysis with selected demographic variables and clinical variables. To find the association between coping strategies among patients undergoing hemodialysis with selected demographic variables and clinical variables.

**Methodology:** A descriptive cross-sectional study was conducted in A.C.S Medical College and Hospital and Government Medical College Hospital, Thiruvallur. The 100 samples who fulfilled the inclusion criteria were selected by using purposive sampling technique. The tool used to collect data were demographic proforma, DASS-21 and brief COPE inventory.

**Findings:** The study findings revealed that among 100 patients undergoing hemodialysis 74(74%) had extreme severe depression, 46(46%) had mild anxiety, 48(48%) had mild stress, 60(60%) had low level of problem-focused coping, 59(59%) had average level of emotion-focused coping, 62(62%) had low level of avoidant coping. However, the overall analysis of coping strategies among patients undergoing hemodialysis revealed that 93(93%) had average coping and 7(7%) had low level of coping. The calculated Karl Pearson's Correlation value of 'r' between depression ( $r = -0.324$ ), anxiety ( $r = -0.348$ ), stress ( $r = -0.303$ ) showed a negligible negative correlation with coping strategies, which was found to be statistically significant at  $p < 0.001$ . The demographic variable occupation ( $\chi^2 = 28.975$ ,  $p = 0.024$ ), marital status ( $\chi^2 = 21.482$ ,  $p = 0.044$ ) had shown statistically significant association with level of depression, anxiety. Respectively occupation ( $\chi^2 = 43.532$ ,  $p = 0.0001$ ) had shown statistically significant association with level of stress. The clinical variable how many years did you receive hemodialysis ( $\chi^2 = 22.358$ ,  $p = 0.034$ ), how many days in a week, did you receive hemodialysis ( $\chi^2 = 20.637$ ,  $p = 0.008$ ), family history of renal problem ( $\chi^2 = 11.405$ ,  $p = 0.022$ ) and access for hemodialysis ( $\chi^2 = 13.211$ ,  $p = 0.010$ ) had shown statistically significant association with level of depression, stress among patients undergoing hemodialysis at  $p < 0.05$  level. The demographic variable marital status ( $\chi^2 = 10.442$ ,  $p = 0.015$ ) had shown statistically significant association with level of coping strategies among patients undergoing hemodialysis at  $p < 0.05$ .

**Conclusion:** The findings of the study showed that the patients undergoing hemodialysis has extreme severe depression, mild level of anxiety and stress. Patient using average level of coping strategies. The study pointed out the importance of reducing the depression, anxiety, stress and improving coping strategies among patients undergoing hemodialysis.

**Keywords:** Depression, anxiety, stress, coping strategies, hemodialysis

### Introduction

"Worry is my worst enemy ... an enemy I unleashed upon myself".

-Terri guillmets.

The urinary system is one of the body's excretory systems. It is made up of two kidneys secrete urine, two ureters transport urine from the kidney to the urinary bladder, and one urinary bladder collects and temporarily stores pee. The urine is evacuated from the urinary bladder to the outside through one urethra. The kidneys contain a pair of excretory

part of the body that eliminate metabolic waste through urine while also maintaining electrolyte and water balance in the body. Chronic kidney disease (CKD) involves progressive, irreversible destruction of the nephrons in both kidneys, the disease process progresses until most nephrons are destroyed and replaced by non-functional scar tissue. According to NIDDK (National Institute of Diabetes and Digestive and Kidney Diseases) hemodialysis is a treatment that filters wastes and water from the blood in the same way as the kidneys did. Hemodialysis can improve the quality of

life and extend our life, but it is not a cure for kidney failure. Hemodialysis (HD) is the most common Renal Replacement Therapy modality in India.

### Background of the study

Individuals diagnosed with kidney disease usually develop psychiatric complications as the dialysis procedure causes changes in the physical health and social life. According to the findings of a study by (Nazemian *et al.*, 2008) <sup>[34]</sup> done a study on association between dialysis patients' depression rates and stress, 64.5% of patients suffer from depression, 51.4% from explicit stress, and 49.7% from hidden stress.

### Need for the study

Dialysis is a stressful process and follows various psychological and social problems which can lead to patients' mental disturbances. Porkodi *et al.*, (2018) <sup>[23]</sup> conducted a cross sectional study to assess the level of stress and coping among 60 Indian patients subjected to hemodialysis at dialysis unit in tertiary care hospital. The result revealed that among patient subjected for dialysis 39 (65%) have mild stress, 12(20%) have moderate stress, 38(63.3%) of the participants never used coping strategies whereas 22(36.7%) of them had sometimes used coping strategies.

### Statement of the problem

A study to assess depression, anxiety, stress and coping strategies among patients undergoing hemodialysis at selected hospitals.

### Objectives

#### Objectives of the study were to

- Assess the level of depression, anxiety, stress among patients undergoing hemodialysis.
- Assess coping strategies among patients undergoing hemodialysis.
- assess correlation between depression, anxiety, stress and coping strategies among patients undergoing hemodialysis.
- Find the association between the depression, anxiety, stress among patients undergoing hemodialysis with selected demographic variables and clinical variables.
- Find the association between coping strategies among patients undergoing hemodialysis with selected demographic variables and clinical variables.

### Hypothesis

H1: There will be a significant correlation between depression, anxiety, stress and coping strategies among patients undergoing hemodialysis.

H2: There will be a significant association between depression, anxiety, stress among patients undergoing hemodialysis with selected demographic variables and clinical variables.

H3: There will be a significant association between coping strategies among patients undergoing hemodialysis with selected demographic variables and clinical variables.

### Materials and Methods

Research approach: Quantitative.

Research design: Descriptive study.

Setting of the study: A.C.S Medical College And Hospital And

Government Medical College Hospital, Thiruvallur

Population: Patients undergoing hemodialysis.

Sample: Patients undergoing hemodialysis who meet the inclusive criteria.

Sample size: 100

Sampling technique: Purposive sampling.

### Sampling criteria

#### Inclusive criteria

1. Patient who are available at the time of data collection.
2. Patient who can able to speak Tamil or English.
3. Patient who is on AV fistula or Internal Jugular Vein access for hemodialysis.

#### Exclusive criteria

1. Patient who are not willing to participate.
2. Patient who are having any cognitive impairment.

### Tools for data collection

#### Section A: Demographic proforma

Demographic proforma includes demographic variables and clinical variables.

#### Section B: Depression, Anxiety, Stress Scale -21

DASS - 21 was designed in 1995 by Lovibond & Lovibond. DASS-21 is a modified version of DASS-42 Self- reported items, each reflecting a negative emotional symptom. These scores ranged from 0- did not apply to me at all (NEVER), 1- applied to me some of the time (SOMETIMES), 2- applied to me to a considerable time (OFTEN), 3- applied to me very much (ALMOST ALWAYS).

#### Section C: Brief COPE Inventory

Brief COPE Inventory was developed by Charles Carver in 1997. The brief COPE inventory is a 28 item self- report questionnaire designed to measure effective & ineffective ways to cope with a stressful life event. The scale can determine someone's primary coping styles with scores on three subscales:

- Problem focused coping.
- Emotional focused coping.
- Avoidant coping.

### Ethical considerations

Ethical clearance was obtained from the Institutional Ethical Committee of A.C.S Medical College and Hospital, Vellappanchavadi. (No. 370 / 2021 / IEC / ACSMCH Dt. 22.10.2021). Informed consent was obtained from the study participants, confidentiality was maintained throughout the study.

### Pilot study

Pilot study was conducted on 05.11.2021 to 06.11.21 with 10 patients undergoing hemodialysis at A.C.S Medical College & Hospital. The data was collected using Demographical proforma, DASS-21 Scale, Brief Cope Inventory on interview method. On the basis of the pilot study, tool was found feasible, and so the investigator was able to proceed to the final study using the same tool.

**Data collection procedure**

The data collection period was one month from 15-11-2021 to 15-12-2021. The written permission was obtained from the Honourable Secretary of A.C.S Medical College and Hospital, Vellappanchavadi, and Dean of Government Medical College Hospital, Thiruvallur. Hemodialysis patients who fulfilled the inclusion criteria were selected by

using purposive sampling technique. The researcher introduced herself to the patient undergoing hemodialysis and developed good rapport with them for their co-operation. The researcher assured the participants for the confidentiality of their response.


**Results and Discussion****Table 1:** Frequency and percentage distribution of demographic variables of the patients undergoing hemodialysis. N=100

Demographic Variables	Frequency	Percentage
<b>Age</b>		
30 - 40 years	15	15.0
41 - 50 years	33	33.0
>50 years	52	52.0
<b>Gender</b>		
Male	66	66.0
Female	34	34.0
<b>Educational status</b>		
Primary school level	54	54.0
Secondary school level	13	13.0
Graduate	9	9.0
Illiterate	24	24.0
<b>Marital status</b>		
Married	71	71.0
Unmarried	9	9.0
Widow	18	18.0
Divorced	2	2.0
<b>Occupation</b>		
Unemployment	72	72.0
Private	5	5.0
Government	3	3.0
Retired	3	3.0
Self-employee	17	17.0
<b>Income</b>		
Below Rs.5000/-	24	24.0
Rs.5,001 - Rs.20,000/-	74	74.0
Above Rs.20,000/-	2	2.0
<b>Residence</b>		
Urban	31	31.0
Rural	69	69.0
<b>Type of family</b>		
Nuclear family	81	81.0
Joint family	19	19.0

**Table 2:** Frequency and percentage distribution of clinical variables of the patients undergoing hemodialysis. N=100

Clinical variables	Frequency	Percentage
<b>History of hereditary disease</b>		
Diabetes mellitus	8	8.0
Hypertension	44	44.0
Both	25	25.0
None	23	23.0
<b>Family history of renal problem</b>		
Yes	16	16.0
No	84	84.0
<b>Dietary habits</b>		
Vegetarian	13	13.0
Mixed	87	87.0
<b>Personal habits</b>		
Smoking	0	0
Alcoholism	0	0
Both a and b	0	0
None	100	100.0
<b>Access for hemodialysis</b>		
AV fistula	96	96.0
Internal Jugular vein	4	4.0
<b>Site of AV fistula</b>		
Right hand	28	28.0


Left hand	72	72.0
<b>How many years did you receive hemodialysis</b>		
Less than 1 year	37	37.0
1 year to 2 years	25	25.0
3 years to 5 years	29	29.0
More than 5 years	9	9.0
<b>How many days in a week did you receive hemodialysis</b>		
2 days or less	84	84.0
3 days	15	15.0
4 days	1	1.0
<b>Do you take medicines regularly</b>		
Yes	79	79.0
No	19	19.0
Sometime	2	2.0



**Fig 1:** Percentage distribution of level of depression, anxiety, stress among patients undergoing hemodialysis

Fig 1 shows that among all the samples 74(74%) had extreme severe depression, 46(46%) had mild anxiety, 48(48%) had mild stress. The above findings were supported by Kumar. V *et al.*, (2018) <sup>[21]</sup> on depression &

anxiety in patients who were undergoing hemodialysis with chronic kidney disease. The study assessed that, 92(61.3%) patients had depression, whereas 42(28%) had anxiety.



**Fig 2:** Percentage distribution of level of coping strategies among patients undergoing hemodialysis



Fig 2 shows that with respect to Problem - focused coping 60(60%) had low level of coping, Emotion - focused coping, 59(59%) had average coping, Avoidant coping 62(62%) had low level of coping. The above findings were supported by Lydia Muthoka *et al.*, (2021) <sup>[29]</sup> the study results concluded that commonly used coping strategies were confrontational (45%), fatalistic (46%) and supportive (48%). The findings were contradictory with the study conducted by Pravan, (2015) the result showed that 53.6% of patients had inadequate coping and 37.9% of patient had moderate coping.

**Table 3:** Correlation between depression, anxiety, stress and coping strategies among patients undergoing hemodialysis. N = 100

Variables	Mean	S.D	Karl Pearson's Correlation 'r' Value
Depression	15.0	3.41	r= -0.324
Coping	66.14	9.23	p=0.001, S***
Anxiety	4.45	1.75	r= -0.348
Coping	66.14	9.23	p=0.0001, S***
Stress	8.79	2.01	r= -0.303 p=0.002, S**

Table 3 shows that even though there was a negligible negative correction between depression, anxiety and stress to coping strategies it provides information to nurses treat patients may not be able to cope appropriately when there is an increase level of depression, anxiety and stress.

#### **Association between depression, anxiety, stress among patients undergoing hemodialysis with selected demographic variables and clinical variables**

The demographic variable occupation, marital status had shown statistically significant association with level of depression, anxiety. Demographic variable occupation had shown statistically significant association with level of stress. The clinical variable years of receiving hemodialysis had shown statistically significant association with level of depression. The clinical variable days in a week receive hemodialysis, family history of renal problem and access for hemodialysis had shown statistically significant association with level of stress. Where as the other clinical variables had shown no significant association with level of stress, anxiety among patients undergoing hemodialysis.

The above findings were supported by Badema Cengi *et al.*, (2010) <sup>[8]</sup> study on depression in hemodialysis patients at Bosnia & Herzegovina. The employment status was found to be significantly associated with depression ( $p < 0.002$ ) whereas unemployed patients were significantly more depressed in relation to employed patients. The above findings were contradictory by Rayyani M. *et al.*, (2014) <sup>[46]</sup> study on stressors and coping strategies in dialysis patient at Iran. The result showed that, there was no significant statistical difference between the type of stressors & demographic variables of occupation ( $p = 0.292$ ) and hemodialysis times per week ( $p = 0.292$ ).

#### **Association between coping strategies among patients undergoing hemodialysis with selected demographic variables and clinical variables**

The demographic variable marital status had shown

statistically significant association with level of coping strategies. However the other demographic variables had shown no significant association with level of coping strategies among patients undergoing hemodialysis. None of the clinical variables had shown significant association with level of coping strategies among patients undergoing hemodialysis.

The above findings were supported by Theodoritsi. *et al.*, (2016) <sup>[47]</sup> The result concluded that social and family support had significant association with marital status  $p < 0.001$ . The above study findings were contradictory by Raayani M. *et al.*, (2014) <sup>[46]</sup> study on stressors and the result founded that, there was no statistically significant difference between the level of coping mechanisms with marital status ( $p = 0.796$ ).

#### **Conclusion**

The findings of the study showed that the patients undergoing hemodialysis has extreme severe depression, mild level of anxiety and stress. Patient using average level of coping strategies. The study pointed out the importance of reducing the depression, anxiety, stress and improving coping strategies among patients undergoing hemodialysis.

#### **Nursing implication**

##### **Nursing practice**

- Nurses help the patients who are undergoing hemodialysis to initiate positive thoughts and motivate them.
- Nursing personnel should help the patients undergoing hemodialysis organizes daily life skills to meet with the challenges of living in a stressful condition.

##### **Nursing education**

- To create awareness and improve the knowledge regarding the psychological problems.

##### **Nursing administration**

- The nursing administrator promote in-service education and special training programs.
- Nursing professionals can offer opportunity to create awareness among nursing students, staffs.

##### **Nursing research**

- There is a need to carry out more research to detect strategies that can be effectively administered to reduce the psychological problems such as depression, anxiety, stress.

#### **Limitations**

- The research was only conducted among patients undergoing hemodialysis.
- The research was done only in two hospitals.

#### **Recommendation for further study**

- Use any intervention for patients undergoing hemodialysis.
- A similar study can be concluded with qualitative approach.

#### **Conflict of Interest**

Not available

## Financial Support

Not available

## Reference

1. Al-Jabi SW, Sous A, Jorf F, Taqatqa M, Allan M, Sawalha L, *et al.* Depression among end-stage renal disease patients undergoing hemodialysis: A cross-sectional study from Palestine. *Renal Replacement Therapy*. 2021;7(1):1-1. <https://doi.org/10.1186/s41100-021-00331-1>
2. Amirkhani M, Shokrpour N, Bazrafcan L, Modreki A, Sheidai S. The effect of resilience training on stress, anxiety, depression, and quality of life of hemodialysis patients: A randomized controlled clinical trial. *Iranian Journal of Psychiatry and Behavioral Sciences*. 2021;15(2). <https://doi.org/10.5812/ijpbs.104490>
3. Awasthi A, Gadia P, Jain S, Koolwal GD. Depression and anxiety in patients of chronic kidney disease undergoing haemodialysis: A study from Western Rajasthan. *Journal of Family Medicine and Primary Care*. 2020;9(8):4282. [https://doi.org/10.4103/jfmpc.jfmpc\\_840\\_20](https://doi.org/10.4103/jfmpc.jfmpc_840_20)
4. Berma A, Abo El-Ata A, Wahba Nadia, Elmwafy R. Relation between stressors, coping strategies and self-efficacy among patients undergoing hemodialysis. *Port Said Scientific Journal of Nursing*. 2021;8(1):122-141. <https://doi.org/10.21608/pssjn.2021.62243.1084>
5. Brito DC, Machado EL, Reis IA, Carmo LP, Cherchiglia ML. Depression and anxiety among patients undergoing dialysis and Kidney Transplantation: A cross-sectional study. *Sao Paulo Medical Journal*. 2019;137(2):137-147. <https://doi.org/10.1590/1516-3180.2018.0272280119>
6. Bharati J, Jha V. Global dialysis perspective: India. *Kidney 360*. 2020;1(10):1141-1145. <https://doi.org/10.34067/kid.0003982020>
7. Bujang MA, Musa R, Liu WJ, Chew TF, Lim CTS, Morad Z. Depression, anxiety and stress among patients with dialysis and the association with quality of life. *Asian Journal of Psychiatry*. 2015;18:49-52. <https://doi.org/10.1016/j.ajp.2015.10.004>
8. Cengic B, Resić H. Depression in hemodialysis patients. *Bosnian Journal of Basic Medical Sciences*. 2010;10(1):73. <https://doi.org/10.17305/bjbms.2010.2653>
9. Chugh KS. Five Decades of Indian nephrology: A personal journey. *American Journal of Kidney Diseases*. 2009;54(4):753-763. <https://doi.org/10.1053/j.ajkd.2009.06.027>
10. Dehkordi AK, Tayebi A, Ebadi A, Sahraei H, Einollahi B. Effects of aromatherapy using the damask rose essential oil on depression, anxiety, and stress in hemodialysis patients: A clinical trial. *Nephro-Urology Monthly*. 2017;9(6). <https://doi.org/10.5812/numonthly.60280>
11. El-Monshed A, Mahgoub N, El-Boraie O, El-Etreby R. Correlation between depression, anxiety and coping strategies among patients on maintenance hemodialysis: A pilot study. *Mansoura Nursing Journal*. 2017;4(1):177-185. <https://doi.org/10.21608/mnj.2017.176420>
12. Gerasimoula K, Lefkothea L, Maria L, Victoria A, Paraskevi T, Maria P. Quality of life in hemodialysis patients. *Materia Socio Medica*. 2015;27(5):305. <https://doi.org/10.5455/msm.2015.27.305-309>
13. Gerogianni G, Polikandrioti M, Babatsikou F, Zyga S, Alikari V, Vasilopoulos G, *et al.* Anxiety-depression of dialysis patients and their caregivers. *Medicina*. 2019;55(5):168. <https://doi.org/10.3390/medicina55050168>
14. Ghasemi Bahraseman Z, Mangolian Shahrabaki P, Nouhi E. The impact of stress management training on stress-related coping strategies and self-efficacy in hemodialysis patients: A randomized controlled clinical trial. *BMC Psychology*. 2021;9(1):1-9. <https://doi.org/10.1186/s40359-021-00678-4>
15. García-Martínez P, Ballester-Arnal R, Gandhi-Morar K, Castro-Calvo J, Gea-Caballero V, Juárez-Vela R, *et al.* Perceived stress in relation to quality of life and resilience in patients with advanced chronic kidney disease undergoing hemodialysis. *International Journal of Environmental Research and Public Health*. 2021;18(2):536. <https://doi.org/10.3390/ijerph18020536>
16. <https://medalerthelp.org/blog/kidney-disease-statistics/>
17. <https://www.niddk.nih.gov/health-information/kidney-disease/kidney-failure/hemodialysis>
18. Joseph N, Thomas A, Philip AS, *et al.* A study to assess the level of stress regarding hemodialysis among chronic renal failure patients in a selected hospital at Ernakulam, Kerala, India. *International Journal of Research and Review*. 2017;4(11):23-26. E-ISSN: 2349-9788; P-ISSN: 2454-2237
19. Jha V, Ur-Rashid H, Agarwal SK, Akhtar SF, Kafle RK, Sheriff R, *et al.* The state of nephrology in South Asia. *Kidney international*. 2019;95(1):31-37. <https://doi.org/10.1016/j.kint.2018.09.001>
20. Khan A, Khan AH, Adnan AS, Sulaiman SA, Mushtaq S. Prevalence and predictors of depression among hemodialysis patients: A prospective follow-up study. *BMC Public Health*. 2019;19(1):1-3. <https://doi.org/10.1186/s12889-019-6796-z>
21. Kumar V, Khandelia V, Garg A. Depression and anxiety in patients with chronic kidney disease undergoing hemodialysis. *Ann Indian Psychiatry [serial online]* 2018 [cited 2022 May 4];2:115-9. Available from: <https://www.anip.co.in/text.asp?2018/2/2/115/246526>
22. Khan A, Khan AH, Adnan AS, Sulaiman SA, Mushtaq S. Prevalence and predictors of depression among hemodialysis patients: A prospective follow-up study. *BMC Public Health*. 2019;19(1). <https://doi.org/10.1186/s12889-019-6796-z>
23. Maria Juliana J, Porkodi Arjunan. stress and coping among Indian Haemodialysis patients *International Journal of Pharmacy and Biological Sciences - ISSN:2321-3272 IJPBS*. 2018, Issue 4| OCT-DEC| 2015 | 08-23
24. Madhyastha S. In *Manipal Manual of Anatomy for Allied Health Science Courses*. Essay, CBS; c2007. p. 289-295.
25. Marthoenis M, Syukri M, Abdullah A, Tandi TM, Putra N, Laura H, *et al.* Quality of life, depression, and anxiety of patients undergoing hemodialysis: Significant role of acceptance of the illness. *The*

- International Journal of Psychiatry in Medicine. 2020;56(1):40-50.  
https://doi.org/10.1177/0091217420913382
26. Mosleh H, Alenezi M, Al johani S, Alsani A, Fairaq G, Bedaiwi R. Prevalence and factors of anxiety and depression in chronic kidney disease patients undergoing hemodialysis: A cross-sectional single-center study in Saudi Arabia. *Cureus*; c2020. https://doi.org/10.7759/cureus.6668
  27. Mosleh H, Alenezi M, Al Johani S, Alsani A, Fairaq G, Bedaiwi R. Prevalence and factors of anxiety and depression in chronic kidney disease patients undergoing hemodialysis: A cross-sectional single-center study in Saudi Arabia. *Cureus*; c2020. https://doi.org/10.7759/cureus.6668
  28. Musa AS, Pevalin DJ, Al Khalaileh MA. Spiritual well-being, depression, and stress among hemodialysis patients in Jordan. *Journal of Holistic Nursing*. 2017;36(4):354-365.  
https://doi.org/10.1177/0898010117736686
  29. Muthoka L, Maina D, Kimani ST. Strategies used by Kenyan patients on maintenance haemodialysis for coping with stress related to intradialytic events. *African Journal of Nephrology*. 2021;24(1). https://doi.org/10.21804/24-1-4529.
  30. Mafi MH, Moghaddam Zeabadi S, Mafi M, Hosseini Golafshani SZ. Relationship between stressors and coping strategies in Iranian patients undergoing hemodialysis. *Jundishapur Journal of Chronic Disease Care*, In Press (In Press); c2018. https://doi.org/10.5812/jjcdc.84508
  31. Meawad Elsayed EB. The effect of Benson's relaxation technique on anxiety, depression and sleep quality of elderly patients undergoing hemodialysis. *International Journal of Nursing Didactics*. 2019;9(2):23-31.
  32. Nagar K, Vaidya A, Patel K. Depression, anxiety and stress among the patient of chronic kidney disease at Nadiad City, a Cross Sectional Survey; c2021. https://doi.org/10.1101/2021.08.01.21261443
  33. Neupane N, Parajuli P, Mehta RS, *et al*. Stressors and coping strategies among the patients undergoing maintenance haemodialysis at B.P. Koirala institute of health sciences. *Int J Health Sci Res*. 2019;9(6):180-192
  34. Nazemian F, Ghafari F, Poorghaznein T. Evaluation of Depression and Anxiety in Hemodialysis Patients. *Medical journal of mashhad university of medical sciences*. 2008;51(3):171-176. Doi: 10.22038/mjms.2008.5511
  35. Najafipour S, Raoofi R, Emamghorieshi F. P-510 - prevalence of depression in hemodialysis patients of hospital in Jahrom Iran. *European Psychiatry*. 2012;27(1). https://doi.org/10.1016/s0924-9338(12)74677-1.
  36. Odette Dorcas TM, Youth TB, Atuhaire C, Priebe G, Cumber SN. Physiological and psychosocial stressors among hemodialysis patients in the Buea Regional Hospital, Cameroon. *Pan African Medical Journal*; c2018. p. 30.  
https://doi.org/10.11604/pamj.2018.30.49.15180
  37. Oneib B, El Filali A, Bentata Y, Ada N. Depression and anxiety disorders in chronic hemodialysis patients and their quality of life: A cross-sectional study about 106 cases in the northeast of Morocco. *Saudi Journal of Kidney Diseases and Transplantation*. 2017;28(2):341. https://doi.org/10.4103/1319-2442.202785
  38. Prema senbulingam K. *Essentials of Medical Physiology* (6<sup>th</sup> ed.). Jaypee Brothers Medical Publishers (P) Ltd; c2012.
  39. R, DC, K, NG, P, D, N, SG, Nagaral JV, N, M, V, RB. Assessment of depression among patients undergoing haemodialysis: A cross-sectional study. *International Journal of Basic & Clinical Pharmacology*. 2019;8(6):1278. https://doi.org/10.18203/2319-2003.ijbcp20192189
  40. Santhi SA, Samson R, Srikanth DP. Effectiveness of physical activity on depression, anxiety, stress and quality of life of patients on Hemodialysis. *Biomedical Research*. 2018;29(9).  
https://doi.org/10.4066/biomedicalresearch.29-18-177
  41. Senmar M, Razaghpour A, Mousavi AS, Zarrinkolah F, Esmaeili F, Rafiei H. Psychological Symptoms in Patients on Dialysis and Their Relationship with Spiritual Well-Being. *Florence Nightingale journal of nursing*. 2020;28(3):243-249.  
https://doi.org/10.5152/FNJJN.2020.19061
  42. Silva RA, Souza VL, Oliveira GJ, Silva BC, Rocha CC, Holanda JR. Coping strategies used by chronic renal failure patients on Hemodialysis. *Escola Anna Nery - Revista De Enfermagem*. 2016;20(1).  
https://doi.org/10.5935/1414-8145.20160020
  43. Sonia AL, Sharma S. (n.d.). In *Medical Surgical Nursing: Volume1 Preparatory Manual for Undergraduates* (1<sup>st</sup> Edition - July 27, 2016; vol 1. essay, Elsevier India.
  44. Ross JSW, WKJ, Waugh A. In *Anatomy and physiology in health and illness*. Essay, Churchill Livingstone, 1996, 338-344.
  45. Zibaei M, Nobahar M, Ghorbani R. Association of stress and anxiety with self-care in hemodialysis patients. *Journal of Renal Injury Prevention*. 2020;9(2):e14-e14.  
https://doi.org/10.34172/jrip.2020.14
  46. Rayyani M, Malekian L, Forouzi MA, Haghdost A, Razban F. Self-care self-efficacy and quality of life among patients receiving hemodialysis in South-East of Iran. *Asian Journal of Nursing Education and Research*. 2014;4(2):165-71.
  47. Theodoritsi A, Aravantinou ME, Gravani V, Bourtsi E, Vasilopoulou C, Theofilou P, Polikandrioti M. Factors associated with the social support of hemodialysis patients. *Iranian journal of public health*. 2016 Oct;45(10):1261.

#### How to Cite This Article

Priyadharshini M, Varghese LS, Hema VH. Depression, anxiety, stress and coping strategies among patients undergoing hemodialysis. *International Journal of Advance Research in Nursing*. 2023;6(1):07-13

#### Creative Commons (CC) License

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International (CC BY-NC-SA 4.0) License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.